



In Vivo Assessment of Antidepressant upshot of *L - Fenchone* in Chronic Unpredictable Mild Stress (CUMS) induced depression like behaviour in Rodents

V. Siva Ganesh*, K. Somasekhara reddy, Raghuvendra varma. P, J. Balaraju.

Ragahavendra Institute of Pharmaceutical Education and Research, Anantapuramu, AP, India.

ABSTRACT

A comprehensive approach like combining the natural medicine and CUS model for evaluating it as anti-depressant would be productive. Fenchone screened for anti-depressant activity in CUMS model using Fluoxetine as standard. Wistar albino rats were selected to CUMS procedure for 28 days and all the period test substance was administered at doses of 400 mg/kg and 800 mg/kg and at the end of the treatment behavioural and biochemical parameters were analyzed and histopathology findings were observed. CUMS exposure caused a depression like behavior corroborated by the increased immobility time in Despair swim test. In Actophotometer decreased locomotor activity and in the hole board test a decrease in the number of head dips and line crossings. Biochemical findings revealed that decreased serum oxidase dismutase and catalase. Fenchone at the doses tested produced significant effects on behavioral and biochemical tests when compared to CUS group. These results manifested that Fenchone had specifically anti-depressant like effect in vivo. In conclusion, the present study advocates that the repeated administration of Fenchone notably reversed CUMS induced depression and oxidative damage and possessed antidepressant like effects, which would be of therapeutic interest for using Fenchone in the treatment of depressive disorders.

Key words:

Chronic Unpredictable Mild Stress,
Despair Swim Test, Catalase,
Super Oxide Dismutase, Fluoxetine.

Article History:

Received On: 29.06.2019,
Revised On: 31.10.2019,
Accepted On: 02.11.2019

Corresponding Author

Name: V. Siva Ganesh
Email: ganesh.cology@gmail.com

1. INTRODUCTION:

Depression is a long-standing malady that influences all age groups. Despite several efficacious antidepressants available today, the ongoing armamentarium of therapy is somewhat sparse, with undesirable results over one in three of all subjects treated¹.

Depression is indicated by disturbances in appetite and sleep, shortfall in cognition and energy. Thoughts of guilt, worthlessness, and suicide are common. Stroke and Coronary artery diseases are more customary in patients having depression, and depression may deliberately aggravate the prediction for patients with a diversity of concomitant aesculapain health².

Depression is distinguished by fierce dejection and desolation, dropping of attentiveness, melancholic worry, deficit pleasure and self-deprecation. Physical changes may occur, significantly in awful depression, including: insomnia, irregular devour patterns, with anorexia and slim down or sporadically overindulging; reduced energy and sex drive; and interference of the normal circadian and ultradian rhythms, body transposition, and multitudinous endocrine functions³.

Even though a maximum number of antidepressants available, 30% to 40% of patients having depression disappoint to acknowledge the first-line antidepressant treatment⁴.

Present day, natural medicines, with their high safety margins, had demonstrated to be productive pharmacotherapy in treating the depression. The model necessitates exposing the rat at unpredictable chronic stress for 28 days to a line of minor-intensity stressors. This outcome in the improvement of an integer of psychological alterations in majority of animals (some animals can be more stress-resistant), including

anhedonia and apathy. This model is considered as good model for antidepressant action⁵.

1.1 HYPOTHESIS OF DEPRESSION⁶

- The receptor sensitivity theory/hypothesis
- Monoamine Theory
- The serotonin only hypothesis
- The latitudinarian hypothesis
- The Electrolyte membrane hypothesis

Stress is a physical rejoinder mechanism but the body has limited capacity to respond⁷

FENCHONE:

SYNONYMS:

1, 3, 3-Trimethyl-2-norbornanone. 1, 3, 3-Trimethyl-2-norcamphanone

SYSTEMATIC NAME:

1, 3, 3-Trimethylbicyclo [2.2.1] heptan-2-one

CHEMICAL CLASS:

Bicyclic monoterpene ketone

Note: many isomers of Fenchone available, and the remaining isomer-specific intimation has been specified: (1R, 4S)-(S)-Fenchone, CAS number: 7787-20-4

(1R, 4S)-(R)-Fenchone, CAS number: 4695-62-9




fig.1: (-)-fenchone

SOURCES: >1.0%

- Lavender (Spanish) 14.9–49.1%; Fennel (bitter) 4.0–24.0%; Thuja 12.2–12.8%; Fennel (sweet) 0.2–8.0%; Cedarwood (Port Orford) 4.7%; Labdanum 1.4–2.3%; Pine (Scots) 0–2.1%

ACUTE TOXICITY:

Oral LD50 in rats 6.16 g/kg and dermal LD50 in rabbits were tested for acute toxicity.

NEUROTOXICITY:

Fenchone was injected subcutaneously, developed clonic epilepsy in mice at 1,133 mg/kg, but not at 500 mg/kg. Given to dogs for 16 days orally, (h)-Fenchone was convulsive and showed death at 1,400 mg/kg/day, produced mild jerking at 750 mg/kg/day, and had no significant effects at 210–420 mg/kg/day.

GENOTOXICITY AND MUTAGENICITY:

Fenchone was not genetically toxic and mutagenic. -Non mutagenic in Salmonella typhi strains, TA98, TA97, TA100 or TA1535, with or without S9 in rat bone marrow micronucleus test. While, Fenchone was not genetically toxic, though at a maximum dose (2,500 mg/kg ip for 3 days), the results were considered equivocal.

SUMMARY:

Fenchone seems to be non-allergenic, non-irritant and non-toxic⁹.

2. AIM AND OBJECTIVE

AIM:

The present investigation is aimed to study the Antidepressant activity of Fenchone in chronic unpredictable stress induced depression like behaviour in rat model.

OBJECTIVE(S) OF THE STUDY:

To evaluate the Anti-depressant activity of Fenchone by estimating the following parameters:

- ✓ Behavioral parameters such as:
 - Spontaneous locomotor activity,
 - Sucrose preference test,
 - Despair swim test,
 - Hole board test
- ✓ Biochemical parameters like:
 - Superoxide dismutase (SOD)
 - Catalase (CAT)

- ✓ Histopathology of brain.

3. METHODOLOGY

3.1. ANIMALS:

Female Rats measure the weights of (200–220 g) were received from the veterinary college, Bangalore, acclimatized for two weeks before experimentation. Animals are placed 12 hour light/ dark cycle under constant humidity (50± 10%) and temperature (22 ±2° C) and animals were under normal pellet diet and water as much. Here all the experiments were performed between Mornings 10:00 to Evening 04:00. The experimental procedures on animals were in consent with the Committee for the Purpose of Control and Supervision of Experiments on Animals, CPCSEA (Approval Number 878/PO/Re/S/05/CPCSEA/003/2017).

3.2. DRUGS:

Fluoxetine was obtained from Glenmark Pharmaceuticals, India. (-)- Fenchone, SOD standard, Glutathione reduced, DTNB, Epinephrine bitartrate and Tris buffer were procured from Sigma-Aldrich, Bangalore, India. Disodium EDTA, Potassium Dihydrogen Phosphate, Sodium Chloride, sucrose, Sodium bicarbonate, Sodium carbonate, Hydrogen peroxide, Trichloroacetic acid, Sodium phosphate, formaldehyde, Normal saline and hydrochloric acid were obtained from different agencies.

3.3. DRUG TREATMENT:

Fluoxetine, a selective serotonin reuptake inhibitor, used as positive control for antidepressant action. Five groups of six animals in each were assigned randomly: Vehicle control, CUMS plus Vehicle, CUMS plus Fenchone (400 mg/kg), CUMS plus Fenchone (800 mg/kg) and CUMS plus Fluoxetine (20 mg/kg). Fenchone and Fluoxetine are administered orally between 9:30 Am and 10:30 Am once per day for four weeks.

3.4. CHRONIC UNPREDICTABLE MILD STRESS MODEL:

Rats were divided into five groups. (1) Non-Stressed, Control group (2) Stressed, CUS group (3) CUS with Low dose of Test (4) CUS with High dose of Test and (5) CUS with Standard treatment, positive control (Fluoxetine 10 mg/ kg). Each group consisting of six animals (n=6) and every group is placed in individual cages. Control group rats were kept undisturbed in their home cages, where as rats in the CUS group were subjected to various stressors for 28 days (four weeks). The stressors include food deprivation, Water deprivation, over night illumination, Slant cage, cold water immersion, foot shock and forced swimming. To avoid monotony, rats were exposed to these stressors at different times on each day. Behavioral tests were begin at day 29 after 24 hours of last foot shock. After behavioral tests, rat brains were isolated for biochemical analysis and for histopathology¹⁵.

Table 1: Protocol for Chronic Unpredictable Stress (CUS).

DAY/ STRESS	Food deprivation	Slant cage	Over- night illumination	Forced swimming	Water deprivation	Cold water immersion	Foot shock
MON	10:00 a.m ↓						
TUE	10:00 a.m	10:00 a.m ↓					
WED		10:00 a.m	06:00 p.m ↓				
THU			06:00 a.m	10:00 a.m ↓ 11:00 a.m			
FRI					10:00 a.m		
SAT					↓ 10:00 a.m	10:00 a.m ↓ 11:00 a.m	
SUN							10:00 a.m ↓ 11:00 a.m

3.5 BEHAVIORAL PARAMETERS:

3.5.1 Spontaneous Loco motor Activity:

3.5.2 Sucrose preference test:

3.5.3 Despair swim test: ²⁸.

3.5.4 Hole Board Test (HBT)

3.6. OXIDATIVE PARAMETERS IN BRAIN TISSUE HOMOGENATE:

3.6.1. SUPEROXIDE DISMUTASE (SOD)

SOD was estimated by the method of Fridovich and Misra (1967).

3.6.2. CATALASE (CAT):

Catalase was estimated by Hugo E. Aebi method, 1974.

3.7. HISTOPATHOLOGY:

3.8. STATISTICAL ANALYSIS

The outcomes are expressed as the mean $\pm$ SEM. Statistical evaluation was carried out by using one way ANOVA followed by Dunnett's Multiple Comparison Test. $P < 0.05$ was considered to be significant.

4. RESULTS AND DISCUSSION

4.1. BEHAVIORAL PARAMETERS


4.1.1. SPONTANEOUS LOCOMOTOR ACTIVITY:

Table 2: Effect of fenchone on Locomotor Activity

S.NO	GROUPS	LOCOMOTOR SCORES
1	NORMAL CONTROL	409.5 $\pm$ 3.722
2	CUS GROUP	259.7 $\pm$ 3.989****
3	CUS+ Fenchone (400 mg/kg)	303.3 $\pm$ 4.745####
4	CUS+ Fenchone (800 mg/kg)	327 $\pm$ 40274####
5	CUS+ Fluoxetine (10 mg/kg)	379 $\pm$ 6.129###

The values expressed are mean $\pm$ SEM, one way ANOVA followed by Dunnett's test. **** $P < 0.0001$ as compared with

the control, #### $P < 0.0001$ as compared with the CUS group, and ### $P < 0.05$ as compared with the CUS group.




Graph 1: Effect of fenchone on locomotor activity in CUS induced rats was recorded. The values expressed are mean $\pm$ SEM, one way ANOVA followed by Dunnett's test. **** $P < 0.0001$ as compared with the control, #### $P < 0.0001$ as compared with the CUS group and ### $P < 0.05$ as compared with the CUS group.

4.1.2. SUCROSE PREFERENCE TEST:

Table 3: Effect of fenchone on Sucrose Preference.

S.NO	GROUPS	%SUCROSE PREFERENCE
1	NORMAL CONTROL	83.85 $\pm$ 0.9839
2	CUS GROUP	39.38 $\pm$ 1.176****
3	CUS+ Fenchone (400 mg/kg)	54.93 $\pm$ 2.414####
4	CUS+ Fenchone (800 mg/kg)	65.86 $\pm$ 1.445####
5	CUS+ Fluoxetine (10 mg/kg)	74.42 $\pm$ 1.631##

The values expressed are mean $\pm$ SEM, one way ANOVA followed by Dunnett's test. ****P < 0.0001 as compared with the control, #### P < 0.0001 as compared with the CUS group and ## P < 0.05 as compared to CUS group.




Graph 2: Effect of fenchone on sucrose preference in CUS induced rats was recorded. The values expressed are mean $\pm$ SEM, one way ANOVA followed by Dunnett's test. ****P < 0.0001 as compared with the control, #### P < 0.0001 as compared with the CUS group and ## P < 0.05 as compared to CUS group.

4.1.3. DESPAIR SWIM TEST:

Table 4: Effect of fenchone on Duration of Immobility.

S.NO	GROUPS	DURATION OF IMMOBILITY (Sec)
1	NORMAL CONTROL	129.3 $\pm$ 5.011
2	CUS GROUP	197.3 $\pm$ 5.408****
3	CUS+ Fenchone (400 mg/kg)	149.8 $\pm$ 2.892####
4	CUS+ Fenchone (800 mg/kg)	180 $\pm$ 2.852####
5	CUS+ Fluoxetine (10 mg/kg)	168 $\pm$ 3.933##

The values expressed are mean $\pm$ SEM, one way ANOVA followed by Dunnett's test. ****P < 0.0001 as compared with the control, #### P < 0.0001 as compared with the CUS group and ## P < 0.05 as compared with the CUS group.




Graph 3: Effect of fenchone on duration of immobility in CUS induced rats was recorded. The values expressed are mean $\pm$ SEM, one way ANOVA followed by Dunnett's test. ****P < 0.0001 as compared with the control, #### P < 0.0001 as compared with the CUS group and ## P < 0.05 as compared with the CUS group.

4.1.4. HOLE BOARD TEST:

Table 5: Effect of fenchone on Hole board test.

S.NO	GROUPS	NUMBER OF HEAD DIPS (Sec)
1	NORMAL CONTROL	29 $\pm$ 1.653
2	CUS GROUP	8.5 $\pm$ 0.8851****
3	CUS+ Fenchone (400 mg/kg)	15 $\pm$ 0.7303####
4	CUS+ Fenchone (800 mg/kg)	18.33 $\pm$ 0.6146####
5	CUS+ Fluoxetine (10 mg/kg)	25.5 $\pm$ 0.9574##

The values expressed are mean $\pm$ SEM, one way ANOVA followed by Dunnett's test. ****P < 0.01 as compared with the control, #### P < 0.0001 as compared with the CUS group and ## P < 0.05 as compared to CUS group.




Graph 4: Effect of fenchone on number of head dips in CUS induced rats was recorded. The values expressed are mean $\pm$ SEM, one way ANOVA followed by Dunnett's test. ****P < 0.01 as compared with the control, #### P < 0.0001 as compared with the CUS group and ## P < 0.05 as compared to CUS group.

4.2.1. SUPEROXIDE DISMUTASE (SOD):

Table 6: Effect of fenchone on Superoxide Dismutase.

S.NO	GROUPS	SOD (U/mg Protein)	% of Control
1	NORMAL CONTROL	1.81 $\pm$ 0.017	100
2	CUS GROUP	0.54 $\pm$ 0.093****	29.38
3	CUS+ Fenchone (400 mg/kg)	1.26 $\pm$ 0.049####	69.61
4	CUS+ Fenchone (800 mg/kg)	1.43 $\pm$ 0.046###	79
5	CUS+ Fluoxetine (10 mg/kg)	1.56 $\pm$ 0.036#	86.18

The values expressed are mean $\pm$ SEM, one way ANOVA followed by Dunnett's test. ****P < 0.01as compared with the control, , #### P < 0.0001 as compared with the CUS group and # P < 0.05 as compared with the CUS group.



Graph 5: Effect of fenchone on SOD activity in CUS induced rats was recorded. The values expressed are mean $\pm$ SEM, one way ANOVA followed by Dunnett's test. ****P < 0.01as compared with the control,, #### P < 0.0001 as compared with the CUS group and # P < 0.05 as compared with the CUS group.


4.2.2. CATALASE (CAT):

Table 7: Effect of fenchone on catalase.

S.NO	GROUPS	Catalase (U/ mg Protein)	% of control
1	NORMAL CONTROL	243.1 $\pm$ 16.64	100
2	CUS GROUP	107.2 $\pm$ 7.546****	44.09
3	CUS+ Fenchone (400 mg/kg)	159.5 $\pm$ 10.07###	65.61
4	CUS+ Fenchone (800 mg/kg)	200.4 $\pm$ 19.48 ^{ns}	82.43
5	CUS+ Fluoxetine (10 mg/kg)	216.1 $\pm$ 8.252 ^{ns}	88.89

The values expressed are mean $\pm$ SEM, one way ANOVA followed by Dunnett's test.


****P < 0.0001 as compared to control, ### P < 0.001 as compared to CUS group and ns- non significant compared to CUS group.




Graph 6: Effect of fenchone on catalase activity in CUS induced rats was recorded. The values expressed are mean $\pm$ SEM, one way ANOVA followed by Dunnett's test.

****P < 0.0001 as compared to control, ### P < 0.0001 as compared with CUS group and ns- non- significant compared to CUS group.


4.3. HISTOPATHOLOGY




A




B



C



D



E

Fig.4.3: A. Control - Morphology of neurons in brain tissue was normal with no evidence of degenerative changes or inflammation.

Fig.4.3: B. Stress - Indicating several patchy neuronal degenerative changes with aggregation extensively dark pinocytic nuclei in neurons.

Fig.4.3: C. Fenchone 400 mg/ kg -Indicating patchy neuronal degenerative changes with dark pinocytic nuclei.

Fig.4.3: D. Fenchone 800 mg/ kg -Indicating mild degenerative changes and inflammation.

Fig.4.3: E. Fluoxetine 10 mg / kg – No significant degenerative changes or inflammation seen.

5. DISCUSSION:

It is normally trusted that Chronic Unpredictable Stress plays an important role in beginning and worsen of depression²⁹. In this view, a model of CUMS induced depression has been initiated to stimulate the pathogenesis of depression in Homo sapiens. A number of investigations proposed that CUMS can

result (in) long-term behavioral disturbances mimic manifestation of depression and that CUMS prompted depression model can be pre-owned for evaluating the effectiveness of antidepressants through behavioral tests like forced swim tests (FST) and sucrose preference Test (SPT). SPT is a measure of misery like behavioral changes, reduced consumption of sugary solutions. The investigations of current study manifest that rats exposed to CUMS procedure guzzle less sucrose solution when compared to normal rats. Chronic stress has been exhibited to dramatically increase the immobility time of rat in forced swim test, an exhibition of behavioral despair³⁰. Unfailingly, CUMS, as observed in the current study, resulted in an increased immobility time in forced swim test in rats, decrease in number of head dips indicates Fenchone produced antidepressant like action in CUMS exposed rats.

6. CONCLUSION:

In conclusion, abiding fenchone treatment during the tack of CUMS was found to relive CUMS induced depression. Whether there is a central action of fenchone that it is critically salient for its anti-depressive effect will be further investigated.

7. LIST OF ABBREVIATIONS:

CAT: Catalase.

CUMS: Chronic Unpredictable Mild Stress.

DST: Despair Swim Test

HBT: Hole Board Test

SSRI: Selective Serotonin Reuptake Inhibitor.

SOD: Super Oxide Dismutase.

8. BIBLIOGRAPHY:

1. Kalappurayil, Tom Mathew, and Joseph Benny Pulinilkku, A Review of Pharmacognostical Studies on *Moringa oleifera* Lam. Flowers Pharmacognosy Journal, 2016; 9(1): 1-7.
2. Bertram G. Katzung, MD, PhD. Basic and clinical pharmacology, 2012; 12: 521-538.
3. Goodman & Gilman's Manual of Pharmacology and Therapeutics, 2008; 1: 278- 288.
4. Haiying Dong, Zhiying Gao et al., β -Asarone reverses Chronic Unpredictable Mild Stress-Induced Depression-Like Behavior and Promotes Hippocampal Neurogenesis in Rats, *Molecules*, 2014; 19: 5634-5649.
5. Gao-Feng Wu, Shuang Ren et al., Antidepressant effect of taurine in chronic unpredictable mild stress induced depressive rats, 2017; 7: 49-54.
6. Debjit bhowmik et.al. Depression symptoms causes, Medications and Therapies, 2012; 1(3): 1-4.
7. The experience of work related stress across occupations, Journal of Managerial Psychology, Vol. 20 No. 2, pp. 178-187.
8. Essentials of medical pharmacology, 2013; 7: 454-455.
9. Robert Tisserand and Rodney et al., constituent profiles, 2014.
10. Rajeshwari Shastry, Aadhyia Sharma, Vinay Sayeli, Ullal Sheetal Dinkar, Screening of Antidepressant Activity of *Punica granatum* in Mice. Pharmacognosy journal, 2017; 9(1): 27-29.

11. Santosh p, venugopl r. et al., Antidepressant activity of methanolic extract of passiflora foetida leaves in mice, 2011; 3(1):5- 9.
12. Ankur jindal n, radhakrishnan mahesh et al., antidepressant-like effect of etazolate, acyclic nucleotide phosphodiesterase 4 inhibitor—an approach using rodent behavioral antidepressant tests battery, 2012; 689: 125-131.
13. Radhakrishnan Mahesh, Shvetank Bhatt et al., Anti-depressant - Like Effect of Novel 5-HT Receptor Antagonist, (4- 3 benzylpiperazin-1-yl) (3 methoxyquinoxalin-2-yl) methanone (6g) in Acute and Chronic Animal Models of Depression, 2011; 49: 619- 626.
14. Ashok Kumar, B.S 1, Lakshman, K et al., Antidepressant Activity of Methanolic Extract of Amaranthus spinosus, 2014; 5 (1):1-7.
15. Yi Chea, Zhu Zhou et al., Chronic unpredictable stress impairs endogenous antioxidant defense in rat brain Neuroscience Letters, 2015; 584: 208–213.
16. Qing-Qiu Mao, Siu-Po Ip et al., Peony glycosides produce antidepressant-like action in mice exposed to chronic unpredictable mild stress: Effects on hypothalamic-pituitary-adrenal function and brain-derived neurotropic factor. Progress in Neuro-Psychopharmacology & Biological Psychiatry, 2009; 33: 1211–1216.
17. Ossowska1. G, Nowak. G et al., Brain monoamine receptors in a chronic unpredictable stress model in rats, 2001; 108 (3): 311–319.
18. Daniela M. Pechlivanova et al., Effect of long-term caffeine administration on depressive-like behavior in rats exposed to chronic unpredictable stress. behavioral pharmacology, 2012; 23: 339–347.
19. Betina Elfving, Pia Høgh Plougmann et al., Differential brain, but not serum VEGF levels in a genetic rat model of depression. Neuroscience Letters, 2010; 474: 13–16.
20. Keskin. I, Guna. Y et al., Effects of Foeniculum vulgare essential oil compounds, fenchone and limonene, on experimental wound healing. Biotechnic & Histochemistry 2017; 1: 1–9.
21. Aydin Him, Hanefi Ozbek et al., Antinociceptive activity of alpha-pinene and fenchone). Pharmacology online, 2008; 3: 363-369.
22. Y Monica Zuzarte, Maria J. Gonçalves et al., Chemical Composition and Antifungal Activity of the Essential Oils of Lavandula pedunculata. Chemistry & biodiversity, 2009; 6: 1-9.
23. Do-hyoung kim, soon-il kim et al., Repellent Activity of Constituents Identified in Foeniculum vulgare Fruit against Aedes aegypti (Diptera: Culicidae). J. Agric. Food Chem, 2002; 50: 6993–6996.
24. Ranjith Kumar Manoharan, Jin-Hyung Lee and Jintae Lee, Antibiofilm and Antihyphal Activities of Cedar Leaf Essential Oil, Camphor, and Fenchone Derivatives against Candida albicans. August 2017; 8: 1-12.
25. Gerhard Vogel. H, Drug Discovery and Evaluation, second edition, 2002.1: 388- 393.
26. Casarotto PC, Andreatini R. Repeated Paroxetine treatment reverses anhedonia induced in rats by chronic mild stress or dexamethasone. Neuro psychopharmacology, 2007; 17:735–742.
27. C. ARON, P. SIMO et al., Evaluation of a rapid technique for detecting minor tranquilizers. Neuropharmacology, 1971; 10: 459-469.
28. Dinesh Dhingra and Amandeep Sharma, Antidepressant-Like Activity of n-Hexane Extract of Nutmeg (Myristica fragrans) Seeds in Mice, J Med Food, 2006; 9 (1): 84–89.
29. Lee AL, OgleWO, Sapolsky RM. Stress and depression: possible links to neuron death in the hippocampus. Bipolar Disord 2002; 4:117–128.
30. Vincent Castagne, Paul Moser et al., Rodent Models of Depression: Forced Swim and Tail Suspension Behavioral Despair Tests in Rats and Mice. Current Protocols in Pharmacology, 2010; 5.8.1- 5.8.14.
31. Zhengchao Xia1, Hongyan Wei2 et al., Chronic unpredicted mild stress-induced depression alter Saxagliptin pharmacokinetics and CYP450 activity in GK rats. Peer journal, 2016; 1611.