World Journal of Current Medical and Pharmaceutical Research
Content available at www.saap.org.in ISSN: 2582-0222

ASSESSMENT OF VISUAL ACUITY AND CLINICAL OUTCOMES IN PATIENTS WITH GLAUCOMA, RETINAL DISEASE AND CATARACT

DURGESH HARI NARAYAN GAUTAM

Assistant Professor, Department of Ophthalmology, Mahayogi Gorakhnath University, Gorakhpur, Uttar Pradesh, India.

DOI: https://doi.org/10.37022/wjcmpr.v8i3.462

ARTICLE HISTORY
Received on: 04-08-2026
Revised on: 19-10-2026
Accepted on: 26-10-2026

Keywords: *Visual acuity; glaucoma; retinal disease; cataract; clinical outcomes; optical coherence tomography; visual field; ophthalmic assessment.*

Corresponding Author
Durgesh Hari Narayan Gautam

ABSTRACT

Background: Glaucoma, retinal diseases and cataract are major causes of visual impairment and blindness worldwide. Although these conditions affect different anatomical structures of the eye, they may produce overlapping symptoms and substantial limitations in daily functioning. Accurate assessment of visual acuity, structural abnormalities and disease-specific visual function is essential for diagnosis, treatment planning and monitoring clinical outcomes.

Objective: To review the methods used to assess visual acuity and visual function in patients with glaucoma, retinal diseases and cataract, and to discuss clinical outcomes associated with current management strategies.

Methods: This narrative review summarizes established ophthalmic guidance and published literature concerning visual acuity assessment, diagnostic investigations, treatment approaches and clinical outcomes in the three disease groups. It focuses on routine clinical assessment and the limitations of visual acuity as a single measure of visual function.

Discussion: Visual acuity provides an important measure of central visual resolution but does not adequately represent every aspect of visual function. In glaucoma, visual-field testing and optic nerve assessment are essential for identifying functional deterioration. Retinal diseases require evaluation of retinal structure and function, often supported by optical coherence tomography and fundus imaging. Cataract assessment includes visual acuity measurement, lens examination and consideration of glare, contrast sensitivity and functional limitations. Treatment outcomes depend on disease severity, timing of intervention, associated ocular conditions and appropriate follow-up.

Conclusion: A comprehensive approach combining visual acuity measurement with disease-specific structural and functional investigations provides a more reliable assessment of ocular health and clinical outcomes than visual acuity alone.

This article is licensed under a Creative Commons Attribution-Non-commercial 4.0 International License.
Copyright © 2026 Author(s) retains the copyright of this article.

INTRODUCTION

Good vision supports everyday activities, including reading, moving around safely, working and maintaining independence. When vision is impaired, these activities can become difficult and may affect a person's social and emotional well-being. Glaucoma, retinal diseases and cataract are common causes of visual loss, but they affect the eye in different ways and therefore require different approaches to assessment and care. Glaucoma refers to a group of diseases in which the optic nerve is progressively damaged. The resulting visual-field defects may develop slowly, and patients may not notice early changes. Central visual acuity can remain good even while peripheral vision is being lost. For this reason, a visual acuity chart alone cannot reliably detect or monitor glaucoma progression; optic nerve examination and visual-field testing are also needed [1,2]. Retinal diseases affect the light-sensitive tissue at the back of the eye and may involve the macula or retinal blood vessels. Examples include diabetic retinopathy, age-related macular degeneration, retinal vein occlusion and retinal detachment. Their effects depend on the site and extent of damage. Some patients develop blurred or distorted central vision, while others experience missing areas of vision or peripheral field loss. Assessment should therefore combine visual-function testing with examination and imaging of the retina [3,4]. Cataract develops when the eye's natural lens becomes cloudy and prevents light from passing through clearly. People may notice blurred vision, troublesome glare, or difficulty seeing in dim light. Surgery can often improve vision when the cataract is responsible for the symptoms, although the result may be limited by other conditions such as glaucoma or retinal disease [5]. These conditions share a common clinical concern: determining the severity of visual impairment and evaluating

whether management preserves or improves useful vision. Visual acuity measurement is central to this process, but its interpretation must be integrated with other examination findings. This review discusses visual acuity assessment, disease-specific investigations, treatment outcomes and the clinical value of comprehensive ophthalmic evaluation.

2. METHODS OF LITERATURE REVIEW

This narrative review discusses published guidance and clinical literature on assessing visual acuity and managing glaucoma, retinal diseases and cataract. The review focuses on visual acuity measurement, visual-field testing, optic nerve examination, retinal imaging, cataract assessment and factors that influence treatment outcomes. Particular attention is given to the ways in which coexisting eye conditions can affect visual recovery.

This is a narrative review, not a registered systematic review. It does not include original patient data or a pooled statistical analysis. The discussion brings together existing evidence and established clinical principles; it should not be interpreted as a new estimate of treatment effectiveness.

3. ASSESSMENT OF VISUAL ACUITY AND VISUAL FUNCTION

3.1 Visual acuity measurement

Visual acuity describes how clearly a person can see fine detail. It is usually measured with a Snellen chart or a standardized logMAR chart. Each eye should generally be tested separately, with suitable refractive correction when needed. Near vision can also be measured when symptoms or the clinical examination make it relevant.

Presenting visual acuity records how well the patient sees with their usual glasses, if any. Best-corrected visual acuity records vision after appropriate refraction. Comparing these results helps clinicians judge whether reduced vision may be due to an uncorrected refractive error or to an underlying eye condition.

For research and clinical follow-up, consistent testing conditions are important. Chart type, testing distance, illumination, correction status and scoring method should be documented whenever possible. LogMAR measurements are particularly useful for quantitative comparisons because they provide a standardized numerical scale.

A standard acuity test mainly measures central vision under high-contrast conditions. It does not fully show how well a person sees to the side, distinguishes objects with similar shades, copes with glare or sees in low light. These gaps are especially important in glaucoma and cataract assessment [1,5].

3.2 Additional measures of visual function

A comprehensive visual assessment may include contrast sensitivity, visual fields, colour vision, glare testing and patient-reported visual functioning. The choice of additional tests depends on the suspected diagnosis and clinical question. Automated perimetry evaluates differential light sensitivity at different locations within the visual field and is especially important in glaucoma, where characteristic defects can occur despite relatively good central visual acuity.

Contrast sensitivity and glare testing can help explain functional difficulties that are not adequately reflected by standard acuity measurements. Patient-reported outcome instruments may document difficulties with reading, mobility, driving and everyday activities. Such assessments complement examination findings, although subjective reports should be interpreted alongside objective findings [2,12].




Figure 01A: Ocular changes in glaucoma, retinal disease and cataract. (A) Optic nerve head cupping in glaucoma. (B) Retinal changes with macular oedema in diabetic retinopathy. (C) Lens opacity in cataract.

4. GLAUCOMA: VISUAL ASSESSMENT AND CLINICAL OUTCOMES

4.1 Pathophysiology and clinical presentation

Glaucoma involves progressive damage to retinal ganglion cells and the optic nerve. Elevated intraocular pressure is an important modifiable risk factor, although glaucoma can develop in patients whose measured pressure remains within the statistically normal range. Disease progression varies according to glaucoma subtype, baseline damage, individual susceptibility and other clinical factors.

Primary open-angle glaucoma often develops gradually and may remain asymptomatic during early stages. Patients may initially retain central visual acuity while experiencing peripheral visual-field loss. Advanced disease can affect central vision and produce substantial functional disability. Angle-closure glaucoma may present differently, including acute symptoms in some cases that require urgent treatment [1,2].

4.2 Clinical assessment

Evaluation of suspected or established glaucoma should extend beyond visual acuity measurement. Important components include intraocular pressure measurement; slit-lamp examination and gonioscopy; optic nerve head assessment and documentation of the cup-to-disc ratio; retinal nerve fibre layer and ganglion cell imaging where available; standard automated perimetry; and assessment of relevant risk factors, previous treatment and disease progression [1,2].

Optical coherence tomography can assist in detecting structural changes in the optic nerve and surrounding retinal nerve fibre layer. Visual-field testing provides complementary functional information. Neither structural imaging nor a single visual-field result should be interpreted in isolation when making management decisions [1,2].

4.3 Management and clinical outcomes

The aim of glaucoma treatment is to protect the optic nerve and preserve the vision that remains. Depending on the type and severity of glaucoma, treatment may include pressure-lowering eye drops, laser treatment or surgery. The choice is guided by disease progression, response to treatment and the patient's circumstances [1].

A clinically meaningful outcome is not necessarily an improvement in visual acuity. In established glaucoma, successful management may consist of maintaining visual acuity, controlling intraocular pressure and preventing or slowing additional visual-field loss. Treatment response

should therefore be assessed using serial pressure measurements, structural examinations and visual-field testing [1,2].

Medication adherence, adverse effects, access to follow-up and timing of diagnosis can influence long-term outcomes. Patients may continue to experience functional limitations even when intraocular pressure is controlled, particularly if substantial optic nerve damage has already occurred. Regular monitoring remains essential because glaucoma-related visual loss is generally irreversible [1,2].

Coexisting cataract presents an additional challenge. Lens opacity may reduce measured visual-field sensitivity and complicate interpretation of glaucoma progression. Cataract surgery can improve visual acuity in appropriate patients, but the extent of recovery depends partly on underlying optic nerve damage [6].

5. Retinal Diseases: Visual Assessment and Clinical Outcomes

5.1 Spectrum of retinal disorders

Retinal disease includes a broad range of vascular, degenerative, inflammatory and mechanical conditions. Diabetic retinopathy can damage retinal vessels and produce haemorrhages, exudation, macular oedema and proliferative changes. Age-related macular degeneration affects central retinal function, while retinal vein occlusion may cause haemorrhage and macular oedema. Retinal detachment can threaten vision through separation of the neurosensory retina from its supporting tissues.

The visual consequences depend on the affected retinal location, disease severity and duration. Macular involvement commonly produces central visual impairment, while peripheral retinal pathology may affect peripheral vision before central acuity deteriorates. Some retinal diseases progress gradually, whereas others require urgent assessment and intervention [7].

5.2 Assessment of visual acuity and retinal structure

Best-corrected visual acuity is an important functional measurement in retinal disease. Nevertheless, it should be interpreted alongside retinal examination and appropriate imaging. Common investigations include dilated fundus examination, fundus photography, optical coherence tomography, fluorescein angiography or other vascular imaging when indicated, and visual-field testing when peripheral retinal function or associated optic nerve disease requires evaluation.

Optical coherence tomography is particularly useful for identifying macular oedema, changes in retinal thickness and structural abnormalities involving the macula. Fundus imaging supports documentation of diabetic retinopathy and other retinal disorders. Investigations should reflect the suspected diagnosis rather than applying every test to every patient [8,9].

5.3 Treatment approaches and outcomes

Management varies substantially according to the retinal disorder. Diabetic retinopathy requires appropriate systemic risk-factor management and ophthalmic surveillance. Selected patients may benefit from intravitreal anti-vascular endothelial growth factor therapy, laser treatment or vitreoretinal surgery. Neovascular age-related macular degeneration is commonly managed with intravitreal therapy, whereas selected retinal detachments require urgent surgical intervention [9,10].

Clinical outcomes may include improvement in visual acuity, stabilization of existing vision, reduction in retinal oedema, resolution of selected complications or prevention of further deterioration. Anatomical improvement on imaging does not always translate into equivalent functional recovery, particularly when irreversible retinal damage has occurred.

The final visual result may be influenced by disease duration, baseline acuity, macular involvement, treatment adherence, recurrence and coexisting ocular conditions. Patients with advanced macular disease may retain substantial visual disability despite treatment. Low-vision rehabilitation, appropriate visual aids and counselling can support functional independence.

6. CATARACT: VISUAL ASSESSMENT AND CLINICAL OUTCOMES

6.1 Clinical characteristics

Cataract is characterized by opacity of the crystalline lens and may develop through ageing, metabolic disease, trauma, medication exposure or other causes. Common symptoms include blurred vision, glare, reduced contrast sensitivity, difficulty reading and impaired vision under challenging lighting conditions.

The severity of symptoms does not always correspond directly to a single visual acuity measurement. Posterior subcapsular cataract, for example, may produce disproportionate glare or functional difficulty. Coexisting refractive error, corneal disease, glaucoma and retinal pathology can further influence presenting visual status [5].

6.2 Preoperative assessment

Assessment includes presenting and best-corrected visual acuity, refraction when appropriate, slit-lamp examination and evaluation of the patient's functional complaints. The type and severity of lens opacity should be documented.

Additional investigations may be required when visual impairment appears disproportionate to the cataract or when coexisting ocular disease is suspected. These may include retinal examination, macular optical coherence tomography, intraocular pressure measurement and visual-field testing where indicated. Biometry is important for calculating the appropriate intraocular lens power before cataract surgery.

Identifying coexisting retinal or optic nerve disease is important because it helps establish realistic expectations for postoperative visual recovery. A patient with visually significant cataract and advanced glaucoma may experience clearer central vision after surgery without recovering visual field already lost to glaucoma [5,6].

6.3 Cataract surgery and postoperative outcomes

For cataract that meaningfully affects daily life, surgery to remove the cloudy lens and replace it with an artificial intraocular lens is the standard treatment. Many patients experience clearer vision and improved ability to carry out everyday tasks, although the result depends on the health of the rest of the eye [5,11].

Postoperative evaluation should document visual acuity, refractive status when appropriate, anterior segment findings and relevant complications. The timing of assessment should be appropriate to the procedure and clinical course. Uncomplicated recovery may differ from recovery complicated by inflammation, corneal oedema,

posterior capsular complications or coexisting retinal disease.

Postoperative visual acuity is influenced by preoperative vision, ocular comorbidities, surgical factors, intraocular lens considerations and postoperative complications. Poor postoperative acuity should prompt assessment for potentially treatable causes, including residual refractive error, macular oedema, corneal abnormalities and posterior segment pathology.

Visual acuity is an important outcome measure but does not capture all benefits of surgery. Reduced glare, improved contrast and greater ease with daily activities may also be clinically meaningful [5,11].

7. COMPARATIVE ASSESSMENT ACROSS THE THREE DISEASE GROUPS

Glaucoma, retinal diseases and cataract require overlapping but distinct approaches to visual assessment. Although visual acuity is relevant in all three groups, it measures only one aspect of visual function.

Table 01: Comparison of key features, assessment and clinical outcomes in glaucoma, retinal diseases and cataract.

Feature	Glaucoma	Retinal diseases	Cataract
Main anatomical site	Optic nerve and retinal ganglion cells	Retina and macula, varying by disease	Crystalline lens
Typical visual impairment	Peripheral field loss; central acuity may initially be preserved	Central or peripheral visual loss depending on pathology	Blurred vision, glare and reduced contrast sensitivity
Key assessment	Visual acuity, intraocular pressure, optic nerve and visual field	Visual acuity, fundus examination and OCT/imaging	Visual acuity, refraction and lens examination
Important additional tests	Automated perimetry and optic nerve imaging	OCT, fundus photography and angiography when indicated	Glare assessment and retinal examination when indicated
Main treatment options	Medication, laser and surgery	Disease-specific treatment, including injections, laser or surgery	Cataract surgery with intraocular lens
Primary treatment objective	Preserve remaining vision and slow progression	Treat retinal disease and preserve or restore function	Improve vision by removing lens opacity
Common clinical outcomes	Stability of visual fields and optic nerve structure	Improvement or stabilization of acuity and retinal status	Improved visual acuity and functional performance

This comparison demonstrates that identical visual acuity measurements may represent different clinical situations. A patient with early glaucoma may have good central acuity but significant peripheral field loss. A patient with macular disease may have central acuity loss despite a relatively clear lens. A patient with cataract may have reduced acuity that improves substantially following surgery, provided no major competing pathology limits recovery. Comparative assessment should therefore consider the anatomical site of disease, the functional deficit and the expected response to treatment.




Figure 02: Comprehensive assessment of visual acuity and clinical outcomes in glaucoma, retinal disease and cataract. The evaluation includes visual acuity measurement, disease-specific assessment, management and follow-up.

8. FACTORS INFLUENCING CLINICAL OUTCOMES

8.1 Disease severity at presentation

Earlier recognition can provide opportunities to intervene before irreversible damage becomes extensive. This is particularly important in glaucoma, where established optic nerve injury cannot generally be reversed. In retinal diseases, delayed treatment may reduce the likelihood of recovering lost visual function. In cataract, surgery can often improve vision, although coexisting pathology may restrict the final outcome [4,13].

8.2 Coexisting ocular disease

Multiple ocular conditions can coexist, particularly among older adults. Cataract may obscure retinal examination or influence visual-field measurements, while glaucoma or macular disease may limit postoperative acuity. Comprehensive assessment is necessary to identify the relative contribution of each condition to visual impairment.

8.3 Treatment adherence and follow-up

Long-term management often depends on adherence to medication, attendance at scheduled examinations and timely access to treatment. This is particularly important for chronic glaucoma and retinal disorders requiring repeated monitoring or intervention. Follow-up permits clinicians to detect progression, assess treatment response and modify management.

8.4 Patient-centred visual functioning

Clinical measurements do not completely represent the effects of visual impairment on everyday life. Reading, mobility, driving, work and self-care may be affected differently across diseases. Patient-reported measures and assessment of functional needs can supplement objective findings and help identify patients who may benefit from rehabilitation or visual aids [12].

9. CHALLENGES AND LIMITATIONS OF VISUAL ACUITY AS AN OUTCOME MEASURE

Visual acuity is reproducible and clinically useful when measured under standardized conditions. However, high-contrast distance acuity may remain normal despite meaningful peripheral field loss. Patients with retinal disease may have metamorphopsia, scotomas or other visual disturbances that are incompletely represented by an acuity score. Cataract may produce glare and reduced contrast sensitivity that interfere with everyday functioning beyond what a standard chart demonstrates.

Visual acuity also requires careful interpretation in longitudinal assessments. Differences between examinations may reflect changes in refractive correction, testing conditions or measurement variability rather than true disease progression. In glaucoma, structural imaging and visual-field results are required to interpret progression. In retinal disease, anatomical changes and disease-specific measures may be necessary. Following cataract surgery, refractive outcomes, complications and patient-reported function may complement acuity measurements [1,2,8].

No single test can describe every aspect of an eye condition. Using visual acuity alongside appropriate imaging, functional tests and follow-up gives a fuller picture of disease status and how well treatment is working [1,2,5,8].

10. CLINICAL IMPLICATIONS AND FUTURE DIRECTIONS

Routine ophthalmic practice should incorporate standardized visual acuity measurement while tailoring additional tests to the suspected or established diagnosis. Consistent documentation supports longitudinal comparisons, clinical communication and evaluation of treatment effectiveness [3,14].

Imaging technologies, automated visual-field analysis and patient-reported outcome instruments provide complementary information. Future clinical research may benefit from standardized definitions of visual improvement, stability and deterioration, as well as clearer reporting of baseline disease severity and follow-up duration.

Comparative research should account for differences between disease groups. Cataract surgery, glaucoma treatment and retinal interventions have different therapeutic objectives, so their outcomes should not be judged solely using the same numerical change in visual acuity. Disease-specific measures should accompany common functional outcomes wherever appropriate.

11. CONCLUSION

Visual acuity assessment is fundamental to the evaluation of patients with glaucoma, retinal diseases and cataract. However, its clinical meaning varies according to the underlying pathology. Glaucoma requires particular attention to optic nerve structure and visual-field progression; retinal diseases require disease-specific retinal examination and imaging; and cataract assessment should include lens opacity, functional complaints and the potential for postoperative recovery.

Clinical outcomes depend on disease severity, timely intervention, associated ocular conditions, treatment adherence and appropriate follow-up. A comprehensive assessment combining visual acuity with structural, functional and patient-reported measures provides a more informative evaluation than visual acuity alone. Individualized management and consistent monitoring remain central to preserving vision and improving visual functioning.

CONFLICT OF INTEREST

The authors declare no conflicts of interest related to this study.

FINDING

Nil

AUTHOR CONTRIBUTION

Both are contributed equally.

ACKNOWLEDGEMENT

Not Declared

REFERENCES

1. Gedde SJ, Bowden EC, Challa P, Vinod K, Kolomeyer NN, Chopra V, Budenz DL. Primary open-angle glaucoma preferred practice pattern. Ophthalmology. 2026;133(4):P1-P103. doi:10.1016/j.ophtha.2025.12.029.
2. American Academy of Ophthalmology. Assessment of visual function in glaucoma: an ophthalmic technology assessment. Ophthalmology. 2011;118(5):986-1002.
3. Wallace DK, Flaxel CJ, Gedde SJ, Jacobs DS, Kopplin LJ, Lee BS, et al. Comprehensive adult medical eye evaluation preferred practice pattern. Ophthalmology. 2026;133(4):P202-P236.
4. World Health Organization. World report on vision. Geneva: World Health Organization; 2019.
5. Miller KM, Oetting TA, Tweeten JP, Carter K, Lee BS, Lin S, et al. Cataract in the adult eye preferred practice pattern. Ophthalmology. 2022;129(1):P1-P126. doi:10.1016/j.ophtha.2021.10.006.
6. Rao HL, Jonnadula GB, Addepalli UK, Senthil S, Garudadri CS. Effect of cataract extraction on visual field index in glaucoma. J Glaucoma. 2013;22(2):164-168.
7. Ferris FL 3rd, Wilkinson CP, Bird A, Chakravarthy U, Chew E, Csaky K, et al. Clinical classification of age-related macular degeneration. Ophthalmology. 2013;120(4):844-851. doi:10.1016/j.ophtha.2012.10.036.
8. Hee MR, Puliafito CA, Wong C, Duker JS, Reichel E, Schuman JS, et al. Optical coherence tomography of the human retina. Arch Ophthalmol. 1995;113(3):325-332. doi:10.1001/archopht.1995.01100030081025.
9. Flaxel CJ, Adelman RA, Bailey ST, Fawzi A, Lim JI, Vemulakonda GA, et al. Diabetic retinopathy preferred practice pattern. Ophthalmology. 2020;127(1):P66-P145. doi:10.1016/j.ophtha.2019.09.025.
10. Early Treatment Diabetic Retinopathy Study Research Group. Photocoagulation for diabetic macular edema: ETDRS report number 1. Arch Ophthalmol. 1985;103(12):1796-1806. doi:10.1001/archopht.1985.01050120030015.
11. Day AC, Donachie PHJ, Sparrow JM, Johnston RL. The Royal College of Ophthalmologists' National Ophthalmology Database study of cataract surgery:

report 1, visual outcomes and complications. Eye (Lond). 2015;29(4):552-560. doi:10.1038/eye.2015.3.

12. Mangione CM, Lee PP, Gutierrez PR, Spritzer K, Berry S, Hays RD. Development of the 25-item National Eye Institute Visual Function Questionnaire. Arch Ophthalmol. 2001;119(7):1050-1058. doi:10.1001/archopht.119.7.1050.
13. Bourne RRA, Steinmetz JD, Flaxman S, Briant PS, Taylor HR, Resnikoff S, et al. Trends in prevalence of blindness and distance and near vision impairment over 30 years: an analysis for the Global Burden of Disease Study. Lancet Glob Health. 2021;9(2):e130-e143. doi:10.1016/S2214-109X(20)30425-3.
14. World Health Organization. Integrated people-centred eye care, including preventable vision impairment and blindness: report by the Director-General. Geneva: World Health Organization; 2019.